arXiv:hep-th/9711033v1 5 Nov 1997

On The Construction of Zero Energy States in Supersymmetric Matrix Models III

Jens Hoppe*
Theoretische Physik, ETH-Hönggerberg
CH-8093 Zürich, Switzerland

Abstract

For a supersymmetric Hamiltonian appearing in the matrix model related to 11 dimensional supermembranes, zero energy states are constructed. A useful symmetry, and an energy-equipartition property is pointed out.

---

* Heisenberg fellow

The supercharges of the previously studied SU(N) matrix model for supermembranes in 11 space-time dimensions are sums of two terms (each being, separately, nilpotent and defining it's own supersymmetric Hamiltonian):

$$\begin{aligned} Q_\beta &= M_a^{(\beta)}\,\lambda_a \,+\, D_a^{(\beta)}\,\partial_{\lambda_a} \\ Q_\beta^\dagger &= D_a^{(\beta)\dagger}\,\lambda_a \,+\, M_a^{(\beta)\dagger}\,\partial_{\lambda_a} \\ & \qquad\qquad \beta \,=\, 1,\ldots,8 \end{aligned} \tag{1}$$

where $\{\lambda_a,\lambda_b\} \,=\, 0 \,=\, \{\partial_{\lambda_a},\partial_{\lambda_b}\}$, $\{\lambda_a,\partial_{\lambda_b}\} \,=\, \delta_{ab}$, and the M_a and D_a , $a \,=\, (\alpha,A) \,=\, (1,1)\ldots(8,N^2-1)$, are first order differential operators (with respect to bosonic variables $(x_{jA},z_A,\overline{z}_A)_{\substack{j=1\ldots7\\ A=1\ldots N^2-1}}$) , given explicitly in terms of purely imaginary, antisymmetric, 8×8, SO(7) Gamma matrices by

$$\begin{aligned} M_{\alpha A}^{(\beta)} &= \delta_{\alpha\beta}\,i\,q_A \,+\, \Gamma^j_{\alpha\beta}\,\frac{\partial}{\partial x_{jA}} \,-\, \frac{1}{2}\,f_{ABC}\,x_{jB}\,x_{kC}\,\Gamma^{jk}_{\alpha\beta} \\ D_{\alpha A}^{(\beta)} &= \delta_{\alpha\beta}\,2\partial_A \,-\, i\,f_{ABC}\,x_{jB}\,\overline{z}_C\,\Gamma^j_{\alpha\beta}\;, \end{aligned} \tag{2}$$

with $q_A \,:=\, \frac{i}{2}\,f_{ABC}\,z_B\overline{z}_C$ (and f_{ABC} being real, totally antisymmetric structure constants of SU(N)).

In [1], solutions of $Q_\beta\chi = 0$, $Q_\beta^\dagger\Psi = 0$, were constructed that are of the form

$$\begin{aligned} \Psi &= (\mathbf{1} - A)^{-1}\;\Psi^{(h)} \\ \chi &= (\mathbf{1} - B)^{-1}\;\chi^{[h]} \end{aligned} \tag{3}$$

with

$$\begin{aligned} A \,=\, \left(I^\dagger\cdot\lambda\right)\left(D^\dagger\cdot\lambda\right)\;,\; B \,&=\, (I\cdot\partial_\lambda)\,(D\cdot\partial_\lambda) \\ I_{\alpha A}^{(\beta)} \,&:=\, i\,\delta_{\alpha\beta}\,\frac{q_A}{q^2}\;, \end{aligned} \tag{4}$$

and

$$\left(M^\dagger\cdot\partial_\lambda\right)\,\Psi^{(h)} \,=\, 0\;,\; (M\cdot\lambda)\;\chi^{[h]} \,=\, 0\;.$$

Here, I will only consider

$$Q_D^{(\beta)} \,:=\, D_a^{(\beta)}\;\partial_{\lambda_a}\;, \tag{5}$$

satisfying $\{Q_D^{(\beta)}\;,\;Q_D^{(\beta')}\} \,=\, 0$, and giving rise to the non-negative supersymmetric quantum-mechanical Hamiltonian $Q_\beta\,Q_\beta^\dagger \,+\, Q_\beta^\dagger\,Q_\beta$, which, in a particular representation for the Γ matrices, reads

$$\begin{aligned} H_D \,=\, &-\; 4\,\partial_A\,\overline{\partial}_A + f_{ABC}\,x_{jB}\,\overline{z}_C\,f_{AB'C'}\,x_{jB'}\,z_{C'} \\ &+\; 2\,f_{AA'E}\,x_{jE}\left(\delta_{\alpha8}\,\delta_{\alpha j'} - \delta_{\alpha j}\,\delta_{\alpha'8}\right)\;\lambda_{\alpha A}\,\partial_{\lambda_{\alpha'A'}} \end{aligned} \tag{6}$$

(for notational simplicity, β is chosen to be 8).

Despite the fact that (6) does not have a genuinely quartic interaction (it is just a simple harmonic oscillator) the existence of zero-modes of (6) is somewhat non-trivial, and its zero-energy eigenfunction(s) show certain features that one may expect from a solution of the more complicated problem (involving M and D).

To determine the spectrum of (6), diagonalize

$$H'_D \,:=\, 2\,f_{AA'E}\,x_{jE}\left(\delta_{\alpha8}\,\delta_{\alpha'j} - \delta_{\alpha'8}\,\delta_{\alpha j}\right)\lambda_{\alpha A}\,\partial_{\lambda_{\alpha'A'}} \tag{7}$$

by noting that if

$$W_{aa'}\, e^{(\lambda)}_{a'} \;=\; \lambda\, e^{(\lambda)}_{a}\ , \tag{8}$$

$$\Psi' \;=\; e^{(\lambda_1)}_{a_1} \dots e^{(\lambda_l)}_{a_l}\ \lambda_{a_1} \dots \lambda_{a_l} \tag{9}$$

will satisfy

$$W_{aa'}\ \lambda_a\ \partial_{\lambda_{a'}}\ \Psi' \;=\; (\lambda_1 + \ldots + \lambda_l)\ \Psi'\ . \tag{10}$$

For the case at hand, (8) reads

$$\begin{aligned} \lambda\, e^{(\lambda)}_{8A} \;&=\; 2\, f_{AA'E}\ x_{jE}\ e^{(\lambda)}_{jA'} \\ \lambda\, e^{(\lambda)}_{jA} \;&=\; -\,2\, f_{AA'E}\ x_{jE}\ e^{(\lambda)}_{8A'}\ , \end{aligned} \tag{11}$$

implying

$$\lambda^2\, e^{(\lambda)}_{8A} \;=\; +4\, S_{AA'}\ e^{(\lambda)}_{8A'}\ , \tag{12}$$

with

$$\begin{aligned} S_{AA'} \;&:=\; f_{ABC}\ x_{jB}\ f_{A'B'C}\ x_{jB'} \\ &=\; -\sum_{j=1}^{7} (Ad\ X_j)^2_{AA'} \end{aligned} \tag{13}$$

being the real symmetric (positive semi-definite, as $\sum_j Tr\left([X_j, Z]^\dagger\ \ [X_j, Z]\right) \geq 0$) matrix whose eigenvalues $s^{(A)}(x) \;=:\; (\omega^{(A)})^2$, $A = 1 \ldots (N^2 - 1)$, determine the frequencies of the harmonic oscillators in (6).

Clearly, the ground state energy of $H_D^{(0)} \;:=\; -\,4\partial_A\, \overline{\partial}_A + \overline{z}_A S_{AA'} z_{A'}$, is

$$E_D^{(0)} \;=\; 2\left(\omega^{(1)} + \ldots + \omega^{(N^2-1)}\right)\ . \tag{14}$$

Due to (11)/(12), $(W_{aa'})$ has eigenvalues $+2\omega_1, \ldots, +2\omega_{N^2-1}, -2\omega_1, \ldots, -2\omega_{N^2-1}$ (plus $6\cdot(N^2 - 1)$ zeroes). Choosing $l \geq N^2 - 1$ (and $\leq 7\cdot(N^2-1)$), as well as $\lambda_A = -2\omega^{(A)}$ for $A = 1 \ldots N^2 - 1$, one sees that the ground state energy of $H_D^{(0)}$ can be compensated by the lowest eigenvalue of H'_D – proving that $H_D\, \Psi = 0$ if

$$\begin{aligned} \Psi \;=\; f(x)\ \ \cdot\ \ & e^{-\frac{1}{2}\ \omega^{(A)}\,(x)\left(\tilde{x}_8^{(A)^2} + \tilde{x}_9^{(A)^2}\right)}\ \ \cdot\ \prod_{A=1}^{N^2-1} \omega^{(A)} \\ \cdot\ \ & e^{(-2\omega^{(1)})}_{a_1}\,(x) \ldots e^{(-2\omega^{(l)})}_{a_l}\,(x)\ \lambda_{a_1} \ldots \lambda_{a_l}\ , \end{aligned} \tag{15}$$

with $\omega^{(l > N^2-1)} \;:=\; 0$, $\tilde{x}_8^{(A)} + i\, \tilde{x}_9^{(A)} \;:=\; R_{AA'}(x) z_{A'}$ $(S \;=\; R^{-1}\, s\, R)$ – and $f(x)$ subject to $\langle \Psi, \Psi \rangle < \infty$ (to get a SU(N) invariant zero energy state, one may average (15) with respect to the SU(N) action on it). One of the interesting features of (15) is its non-trivial dependence on x, in particular when two eigenvalues cross – or any of the $\omega^{(A)}(x)$ reach(es) zero. It seems that the (almost trivial) supersymmetric harmonic oscillator (6) carries quite a lot of information.

Finally, I would like to mention the following two properties of (supersymmetric) Hamiltonians with supercharges of the type (1), (2):

I) Let N be odd, $\hat{x}_\mu \;:=\; \delta_{\mu 8}\, x_\mu - (1 - \delta_{\mu 8}) x_\mu$, $\mu = 1 \ldots 9$, and

$$* \;:=\; \sum_m (-)^m\ \frac{\epsilon_{a_1 \ldots a_\Lambda}}{m!\,(\Lambda - m)!}\ \lambda_{a_1} \ldots \lambda_{a_m}\ \partial_{\lambda_{a_\Lambda}} \ldots \partial_{\lambda_{a_{m+1}}}\ , \tag{16}$$

i.e.,

$$\begin{aligned} *\ \lambda_a\ * &= \partial_{\lambda_a} \\ *\ \partial_{\lambda_a}\ * &= -\lambda_a\ . \end{aligned} \tag{17}$$

Then

$$*\ Q(\hat{x})\ * \ =\ Q^\dagger\,(x)\ , \tag{18}$$

implying

$$*\ H(\hat{x})\ * \ =\ H\ . \tag{19}$$

If, therefore, a zero-energy state of H exists, it may be assumed to be self-dual,

$$*\ \Psi(\hat{x})\ =\ \Psi(x)\ , \tag{20}$$

which means that in looking for such a state it is sufficient to only solve $Q\,\Psi = 0$ (as $Q^\dagger\Psi = 0$ then follows, via (20), (18)).

II) Let $H = -\vec{\nabla}^{\,2} + V(\vec{x}) + H_F(\vec{x})$, with $V(\lambda\vec{x}) = \lambda^{2n}V(\vec{x})$, $H_F(\lambda\vec{x}) = \lambda^{n-1}H_F(\vec{x})$. $H\Psi = 0$ then implies

$$\langle -\triangle \rangle_\Psi\ =\ \langle V \rangle_\Psi\ , \tag{21}$$

as

$$\begin{aligned} 0 &= \langle \Psi,\ \Big[\vec{x}\cdot\vec{\nabla}, -\triangle + V + H_F\Big]\ \Psi\rangle \\ &= \langle \Psi,\ (2\triangle + 2nV + (n-1)\ H_F)\ \Psi\rangle \\ &= (n+1)\ \langle \Psi,\ (\triangle + V)\ \Psi\rangle\ . \end{aligned} \tag{22}$$

In the case of (1), more detailed, local, relations can be obtained.

Acknowledgement
I would like to thank M. Bordemann, F. Finster, J. Fröhlich, G.M. Graf, L. Motl, R. Suter and S.-T. Yau for valuable discussions.

References:

[1] J. Hoppe; hep-th/9709217.